

# Energy dependence of NBO in Scorpius X-1

J. Wang<sup>1</sup>, H.-K. Chang<sup>1,2</sup>, and C.-Y. Liu<sup>2,3</sup>

<sup>1</sup> Institute of Astronomy, National Tsing Hua University, Hsinchu 30013, Taiwan  
e-mail: jwang@mx.nthu.edu.tw

<sup>2</sup> Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan

<sup>3</sup> LESIA, Paris Observatory, 92195 Meudon, France

the date of receipt and acceptance should be inserted later

**Abstract.** We report the energy dependence of normal branch oscillations (NBOs) in Scorpius X-1, a low-mass X-ray binary (LMXB) Z-source. Three characteristic quantities (centroid frequency, quality factor and fractional root-mean-squared (rms) amplitude) of a QPO signal as functions of photon energy are investigated. We found that the NBO centroid frequency decreases with increasing photon energy when it is below 6-8 keV and turns positively correlated with photon energy at the higher-energy side. The rms amplitude increases significantly with the photon energy below 13 keV, and then flats out in the energy band of 13-20 keV. There is no clear dependence on photon energy for the quality factor. Based on these results, we suggest that the NBO originates mainly from the transition layer.

**Key words.** accretion, accretion disk – stars: individual (Scorpius X-1) – stars: neutron – X-rays: stars

## 1. Introduction

The bright persistent neutron star low-mass X-ray binary (LMXB) Scorpius X-1 traces a "Z" track in the X-ray color-color diagram (CCD), which consists of three branches — the horizontal branch (HB), the normal branch (NB), and the flaring branch (FB, Hasinger & van der Klis 1989). The typical Fourier power spectra in Scorpius X-1 are composed of distinguishably aperiodic variabilities in most of these branches, including noise components (broad structure) and quasi-periodic oscillations (QPOs, narrow feature). There are three distinct types of QPOs observed by Rossi X-Ray Timing Explorer (RXTE), i.e., the normal branch oscillation (NBO) at  $\sim 6$  Hz, the horizontal branch oscillation (HBO) at  $\sim 45$  Hz with a harmonic at about 90 Hz, and the kilohertz (kHz) QPOs (van der Klis et al. 1996, 1997).

The NBOs in Scorpius X-1, which occur on the mid- and lower NB, show some correlations with the variability of the  $\sim 45$  Hz HBO and the twin kHz QPOs, (van der Klis et al. 1996; Yu 2007), and therefore imply some kind of coupling between the three types of QPOs (van der Klis et al. 1996; Dieters & van der Klis 2000). The NBO frequency remains approximately constant at about 6 Hz, in general. However, the timing properties along the Z track shows that the NBO frequency extends to at most  $\sim 21$  Hz and its position moves smoothly into the lower part of the FB (Priedhorsky et al. 1986; Dieters & van der Klis 2000). This smooth transition indicates that the NBO and FBO are physically related to each other (Casella et al. 2006). Properties of both the twin kHz QPO and 45 Hz

HBO depend on the NBO flux. The upper kHz QPO frequency is anticorrelated with the NBO flux, and the lower kHz QPO becomes stronger when the NBO flux is low (Yu et al. 2001). Significant HBOs are detected during the NBO phase of high flux, while the HBO disappears at the low flux of NBO phase (Yu 2007). The coupling between the properties of kHz QPO and HBOs and the phase of the NBO makes the NBO a unique phenomenon.

The photons from different regions of the accretion disk carry different energies and present distinct physical properties. So the QPOs detected in different energy band may have varying characteristics. Inspired by the idea that the photon energy dependence of QPOs may provide some additional information for the nature of QPOs, and limited by the available data which provide enough spectral information and at the same time include QPO signals, we perform a systematically investigation for the energy dependence of NBOs in this work. We describe our data reduction and analysis in section 2. Results are present in section 3. In section 4, we discuss physical implications of our result. Section 5 contains a short summary.

## 2. Data reduction and analysis

To investigate the energy dependence of QPOs in Scorpius X-1, high time resolution data with enough spectral information are needed. QPO signals should also be present in these data. We searched all the RXTE observations of Scorpius X-1 for suitable data. Among all the RXTE Sco X-1 data, two binned data modes, i.e., B\_4ms\_16A\_0\_249\_H and B\_4ms\_16B\_0\_249\_H,

which have a 4-ms time resolution and 16 energy bands composed of all the PCA channels (from original channel 0 to 249) binned with two different schemes (A and B), provide adequate timing and spectral information for our purpose. We examined all the data recorded in those two modes and selected those having QPO signals present (with quality factor  $Q$  larger than 2) in the energy band of channel 0-63, which roughly corresponds to 2 to 20 keV. Those selected data are listed in Table 1. We further defined 6 energy bands, as shown in Table 2, for the energy dependence study. An HBO signal was found in the highest energy band defined in Table 2 in the data set 30035-01-07-00, but cannot be detected in the original 0-63 channel. In addition, another  $\sim 45$  Hz HBO signal was detected in the original 0-63 channel in 93082-01-02-05, but cannot be found in any individual energy band defined in Table 2. We also found another data set, 30035-01-01-00, having the  $\sim 45$  Hz HBO signal in the original 0-63 channels but without NBO. In further examination on this data set, its HBO signal was only detected in the original 0-16 channels but not in higher energy bands. We therefore did not include this data set in this work. Limited by the data available, we study only NBOs in this work. Some of those data sets include several segments. We only exploited the segments which show the NBO signal (see Table 1 for details).

The data were reduced with the HEASOFT package version 6.8. Only those data with the elevation angle  $> 10^\circ$  and the time since South Atlantic Anomaly (SAA)  $< 0$  or  $> 30$  min were selected for further analysis. We extracted light curves with a time resolution of 4 ms ( $2^{-8}$  s) in the six energy bands defined in Table 2. Using the Leahy normalization (Leahy et al. 1983), we computed the PDS without subtracting white noise. The average power expected from a Poisson distribution is 2. The original 4 ms ( $2^{-8}$  s) time resolution was used, and each continuous set was divided into segments of 8192 bins. A fast Fourier transform (FFT) algorithm for each segment was computed yielding the frequency PDS in the range of  $3 \times 10^{-2} - 128$  Hz. We calculated the average PDS for each continuous data interval. A logarithmic rebinning of -1.02 was used for the rebin option. Figure 1 shows an example of the  $\sim 6$  Hz NBO signal.

We employed three components to fit the resulting PDS: a Lorentzian for the very low frequency noise (VLFN), a power law for the white noise, and another Lorentzian for the NBO. All the fits were performed using the  $\chi^2$  minimization technique. The errors of the analytic model parameters were obtained at the  $\chi^2$  variation of 2.70, which corresponds to a 90% confidence for a single parameter.

A QPO signal is characterized by three quantities: the centroid frequency (representing the position of a QPO), the quality factor (characterizing the coherence of a QPO signal), and the fractional root-mean-squared (rms) amplitude (a measure of the signal strength, which is proportional to the square root of the peak power contribution to the PDS). From the fitting we obtained three parameters of NBOs, i.e., the oscillation amplitude, the centroid frequency, and the full width at half maximum (FWHM). The quality factor was then computed according to its definition (centroid frequency/FWHM). The rms amplitude was calculated by the integral of the normalized Lorentzian PDS in the appropriate frequency range and taking into account the mean count rate of the source. Details of rms

**Table 2.** RXTE/PCA energy bands used in this work.

| The observations in 1996 and 1998 |                   |                       |
|-----------------------------------|-------------------|-----------------------|
| PCA Channel                       | Energy Band (keV) | Centroid Energy (keV) |
| 0-13                              | 1.83-5.07         | 2.54                  |
| 14-19                             | 5.07-7.24         | 6.16                  |
| 20-22                             | 7.24-8.34         | 7.79                  |
| 23-28                             | 8.34-10.52        | 9.43                  |
| 29-35                             | 10.52-13.09       | 11.81                 |
| 36-63                             | 13.09-23.46       | 18.26                 |
| The observations in 2009          |                   |                       |
| PCA Channel                       | Energy Band (keV) | Centroid Energy (keV) |
| 0-11                              | 2.18-5.26         | 2.63                  |
| 12-15                             | 5.26-7.02         | 6.14                  |
| 16-19                             | 7.02-8.78         | 7.90                  |
| 20-22                             | 8.78-10.10        | 9.44                  |
| 23-30                             | 10.10-13.60       | 11.85                 |
| 31-63                             | 13.60-28.40       | 21.00                 |




**Fig. 1.** One example of PDS which shows a  $\sim 6$  Hz NBO. The PDS is produced in the PCA original channels 0-63. The mean count rate is 14348 cts/s/PCU. The exposure time is 938 s.

determination can be found in Lewin et al. (1988) and van der Klis (1989). The errors of the quality factor and rms amplitude were estimated with standard error propagation (Bevington & Robinson 2003).

### 3. Results

We investigated the photon energy dependence of three characteristic quantities (i.e. the centroid frequency, quality factor, and rms amplitude) of NBOs. Figure 2 shows the energy dependence of the centroid frequency. Although error bars are large, there seems to be a common trait in almost all the data sets that the dependence evolves from a negative correlation to a positive one with increasing photon energy. We therefore adopt a two-power-law function,  $\nu(E) = a E^{\Gamma_1} + b E^{\Gamma_2}$ , to describe the relation between the centroid frequency  $\nu$  (in units of Hz) and the photon energy  $E$  (in units of keV). The fitting results are listed in Table 3. In all the data set we have a negative

**Table 1.** Data log. Here listed are the RXTE observation identification number (ObsID), their start and stop time, the exposure time in seconds of the light curve after screening (on elevation and time\_since\_SAA) and selection of NBO signal presence, the number of working PCU, the original 0-63 channels count rate in units of cts/s/PCU and the segments with NBO signals.

| ObsID           | Start             | End               | Exposure<br>(s) | PCU | Rate<br>(cts/s/PCU) | Seg |
|-----------------|-------------------|-------------------|-----------------|-----|---------------------|-----|
| 10056-01-04-00  | 27/05/96 20:42:05 | 27/05/96 23:02:13 | 2633            | 5   | 18776               | All |
| 10056-01-04-02  | 28/05/96 04:26:25 | 28/05/96 10:29:13 | 10500           | 5   | 17685               | All |
| 10056-01-05-00  | 28/05/96 11:11:04 | 28/05/96 14:23:13 | 5277            | 4   | 16469               | All |
|                 |                   |                   | 375             | 5   |                     |     |
| 30035-01-02-000 | 31/05/98 00:26:47 | 31/05/98 08:26:47 | 77              | 3   | 18182               | 4,9 |
|                 |                   |                   | 2936            | 5   |                     |     |
| 30035-01-05-00  | 01/06/98 00:26:55 | 01/06/98 01:18:14 | 656             | 3   | 18413               | All |
| 30035-01-07-00  | 02/07/98 06:26:32 | 02/07/98 10:27:14 | 4968            | 5   | 16486               | 1,2 |
| 93082-01-01-01  | 16/06/09 23:43:14 | 17/06/09 00:56:18 | 48              | 4   | 14718               | All |
|                 |                   |                   | 926             | 5   |                     |     |
| 93082-01-01-03  | 18/06/09 01:40:22 | 18/06/09 02:21:18 | 938             | 5   | 14348               | All |
| 93082-01-01-04  | 17/06/09 05:13:12 | 17/06/09 06:01:18 | 1556            | 5   | 14486               | All |
| 93082-01-02-05  | 23/06/09 04:07:09 | 23/06/09 04:40:18 | 1311            | 3   | 13447               | All |

and a positive power index, which together make the centroid frequency decrease with increasing energy first and then when beyond about 6-8 keV it turns to be positively correlated with the photon energy. This dependence is not obvious in one data set (ObsID 30035-01-05-00) among the ten under study. While the small  $\chi^2_\nu$  value indicates that this description is only suggestive due to the large error bars, the appearance of this trait in almost all the data sets asserts its reality. Figure 3 shows the energy dependence of the quality factor. We do not see any clear relation between the quality factor and the photon energy.

In Figure 4 we can see that the rms amplitude increases monotonically with the photon energy except for the highest energy band. It means that the strength of the NBO signal increases with the photon energy. The rms amplitude increases significantly with the photon energy below 13 keV. However, the increase of rms amplitude with photon energy seems to stop when the photon energy is higher than  $\sim 13$  keV, and there might be a plateau. We do not have enough photons at even higher energies for our study. It would be interesting to know beyond which energy the rms amplitude starts to decrease. To describe the increase of the rms amplitude with photon energy, we use a power law relation,  $\text{rms}(E) = C E^\Gamma$ , to fit the data points in the lower five energy bands. Table 4 shows the fitting results. Roughly speaking, the rms amplitude increases with photon energy according to a power law of power index about 0.7 before reaching a possible plateau. In the data set 30035-01-07-00 the plateau seems to start at a lower energy.

#### 4. Discussions

The mechanism of NBOs was suggested to be related to the radial oscillation during radiation feedback (Fortner et al. 1989; Miller & Park 1995) or the different modes of disk oscillations (Alpar et al. 1992; Titarchuk et al. 2001; Bildsten & Cutler 1995; Bildsten et al. 1996; Horak et al. 2004). The standard explanation attributes the  $\sim 6$  Hz NBO to part of the accretion, with near Eddington accretion rate, in an approximately spher-

ically symmetric radial inflow at a radius of  $\sim 100$  km (van der Klis 1995; Hasinger et al. 1990). According to the radiation hydrodynamic model (Fortner et al. 1989), NBO originates from a radiation-force/opacity feedback loop within a spherical flow region at about 300 km from the neutron star surface. The frequency of the oscillations depends on the luminosity. Alternatives are the acoustic oscillation in a thick disk due to the density and optical-thickness perturbations caused by the rotating medium in a subsonic region of the accretion disk (the frequency is determined by the rotation rate and vorticity, Alpar et al. 1992), g-mode oscillations arising from the thermal buoyancy in the ocean on rotating neutron stars (Bildsten & Cutler 1995; Bildsten et al. 1996), the low-frequency modulation in a non-linear resonance oscillation of the relativistic disk (Horák et al. 2004), and the acoustic oscillations of a spherical shell surface within  $\sim 20$  km of the neutron star surface because of the change of accretion geometry in the transition zone at high accretion rate (Titarchuk et al. 2001). All of these interpretations focus on the  $\sim 6$  Hz frequency and its dependence on the accretion rate. None of them can explain the energy dependence of the centroid frequency and rms amplitude.

In the NBOs of Scorpius X-1, both the centroid frequency and rms amplitude display strong energy dependence. The centroid frequency changes from anticorrelation to a positive correlation at about 6-8 keV. Such a nonmonotonic energy dependence of the centroid frequency implies that the emission zone of the NBOs experience a radial variation during the accretion process. The rms amplitude increases monotonically and significantly with the photon energy below  $\sim 13$  keV. When the photon energy is higher than 13 keV, the rms amplitude seems to stop increasing. It may indicate that the NBO signals reach the strongest strength at 13-20 keV. Higher energy photons are often emitted from the inner region of accretion disk. Consequently, the most possible region responsible for such physics can be referred to the transition zone between the inner boundary of accretion disk and the magnetosphere, in which the transition of the properties for accretion flow occurs

**Table 3.** Fitting results of a two-power-law description ( $\nu(E) = a E^{\Gamma_1} + b E^{\Gamma_2}$ ) for the energy dependence of the centroid frequency of NBOs in Scorpius X-1.

| ObsID           | $\Gamma_1$  | a         | $\Gamma_2$  | b         | $\chi^2_\nu$ |
|-----------------|-------------|-----------|-------------|-----------|--------------|
| 10056-01-04-00  | -0.17±0.058 | 3.40±0.65 | 0.11±0.028  | 3.05±0.39 | 0.19         |
| 10056-01-04-02  | -0.22±0.026 | 3.65±0.29 | 0.15±0.030  | 2.79±0.43 | 0.13         |
| 10056-01-05-00  | -0.62±0.023 | 2.52±0.30 | 0.086±0.025 | 5.03±0.38 | 1.03         |
| 30035-01-02-000 | -0.42±0.038 | 2.32±0.43 | 0.083±0.022 | 4.38±0.29 | 0.02         |
| 30035-01-05-00  | -0.43±0.11  | 0.81±1.13 | 0.040±0.054 | 5.23±0.77 | 0.65         |
| 30035-01-07-00  | -3.58±0.27  | 4.27±0.30 | 0.016±0.014 | 6.22±0.20 | 0.37         |
| 93082-01-01-01  | -0.24±0.052 | 5.87±0.70 | 0.35±0.050  | 1.38±0.68 | 0.36         |
| 93082-01-01-03  | -0.16±0.084 | 3.16±1.00 | 0.082±0.029 | 3.65±0.46 | 0.15         |
| 93082-01-01-04  | -0.41±0.057 | 3.52±0.85 | 0.10±0.042  | 4.19±0.68 | 0.62         |
| 93082-01-02-05  | -0.29±0.067 | 3.11±0.78 | 0.11±0.025  | 3.56±0.35 | 0.29         |

**Table 4.** Fitting results of a power law ( $\text{rms}(E) = C E^\Gamma$ ) for the increase of rms amplitude of NBOs in Scorpius X-1. Only the five lower energy bands are included in the fitting.

| ObsID           | $\Gamma$  | C         | $\chi^2_\nu$ |
|-----------------|-----------|-----------|--------------|
| 10056-01-04-00  | 0.78±0.18 | 1.06±0.40 | 0.27         |
| 10056-01-04-02  | 0.71±0.09 | 1.92±0.34 | 0.44         |
| 10056-01-05-00  | 0.75±0.06 | 1.77±0.22 | 0.76         |
| 30035-01-02-000 | 0.73±0.12 | 1.27±0.30 | 0.30         |
| 30035-01-05-00  | 0.63±0.34 | 1.12±0.80 | 0.14         |
| 30035-01-07-00  | 0.70±0.07 | 1.79±0.25 | 3.13         |
| 93082-01-01-01  | 0.70±0.19 | 1.52±0.61 | 0.36         |
| 93082-01-01-03  | 0.74±0.20 | 1.52±0.66 | 0.28         |
| 93082-01-01-04  | 0.72±0.18 | 1.44±0.56 | 0.18         |
| 93082-01-02-05  | 0.70±0.16 | 1.65±0.55 | 0.24         |

(Titarchuk et al. 1998; Titarchuk & Osherovich 1999; Titarchuk et al. 1999). NBO occurs on high accretion rates, i.e. near the Eddington rate. The high mass flux makes more and more matter deposit in the transition zone, leading to the expansion and the radial scale change of this region. However, the pile of more deposit with greater viscosity can contribute to the radiation enhancement, as well as the emission of photons with higher energies. As a result, the radial scale change of the transition zone may be responsible for the observed nonmonotonic energy dependence of the centroid frequency. In addition, because of the limited region of the transition zone, more deposit enhances the viscosity and thus suppresses the further monotonically increasing strength of NBO signal, which can explain the energy dependence of rms amplitude at high energy. We therefore suggest that the NBOs be one kind of oscillations in the transition zone.

## 5. Summary

We investigated the energy dependence of three characteristic quantities of NBOs in Scorpius X-1. We have the following conclusion:

(1). The energy dependence of centroid frequency changes from a negative correlation to a positive one. The turning points are located at 6-8 keV.

(2). It seems that the quality factors have no obvious relation with the photon energy, which indicates that the coherence of NBOs is independent on the photon energy.

(3). The rms amplitude increases significantly with the photon energy below 13 keV. It seems to reach a plateau beyond that.

(4). The observed energy dependence of three characteristic quantities of NBOs in Scorpius X-1 implies that the NBOs originate from one kind of oscillations in the transition layer.

(5). HBO signals are present in three data sets (30035-01-07-00, 93082-01-02-05 and 30035-01-01-00), but they display distinct energy dependence. The HBO signal is only detected in the original 36-63 channels for 30035-01-07-00 and in the original 0-16 channels for 30035-01-01-00. For 93082-01-02-05, the HBO is detected in the original 0-63 channels, but not in any individual energy band defined in Table 2. It seems that the energy dependence of HBO signals changes with the state of the source in Scorpius X-1. Further investigation on this phenomenon will be very rewarding but it requires data of suitable timing and spectral resolution and of good enough statistics.

## 6. Acknowledgments

This work was supported by the National Science Council of the Republic China under grant NSC 99-2112-M-007 -017 -MY3.




**Fig. 2.** The centroid frequency versus the photon energy for NBOs in Scorpius X-1. The solid lines are the best fits with a two-power-law function to the data.

## References

- Alpar, M. A., Hasinger, G., Shaham, J., & Yancopoulos, S. 1992, *A&A*, 257, 627
- Bevington, P. R., & Robinson, D. K. 2003, *Data reduction and error analysis for the physical sciences*, 3rd ed., by Philip R. Bevington, and Keith D. Robinson. Boston, MA: McGraw-Hill, ISBN 0-07-247227-8, 2003
- Bildsten, L., & Cutler, C. 1995, *ApJ*, 449, 800
- Bildsten, L., Ushomirsky, G., & Cutler, C. 1996, *ApJ*, 460, 827
- Casella, P., Belloni, T., & Stella, L. 2006, *A&A*, 446, 579
- Dieters, S. W., & van der Klis, M. 2000, *MNRAS*, 311, 201
- Fortner, B., Lamb, F. K., & Miller, G. S. 1989, *Nature*, 342, 775
- Hasinger, G., & van der Klis, M. 1989, *A&A*, 225, 79
- Hasinger, G., van der Klis, M., Ebisawa, K., Dotani, T., & Mitsuda, K. 1990, *A&A*, 235, 131




**Fig. 3.** The quality factor versus the photon energy for NBOs in Scorpius X-1.

- Horák, J., Abramowicz, M. A., Karas, V., & Kluzniak, W. 2004, *PASJ*, 56, 819
- Leahy, D. A., Darbro, W., Elsner, R. F., et al. 1983, *ApJ*, 266, 160
- Lewin, W. H. G., van Paradijs, J., & van der Klis, M. 1988, *Space Sci. Rev.*, 46, 273
- Miller, G. S., & Lamb, F. K. 1992, *ApJ*, 388, 541
- Miller, G. S., & Park, M.-G. 1995, *ApJ*, 440, 771
- Priedhorsky, W., Hasinger, G., Lewin, W. H. G., et al. 1986, *ApJ*, 306, L91
- Titarchuk, L. G., Bradshaw, C. F., Geldzahler, B. J., & Fomalont, E. B. 2001, *ApJ*, 555, L45
- Titarchuk, L., Lapidus, I., & Muslimov, A. 1998, *ApJ*, 499, 315
- Titarchuk, L., & Osherovich, V. 1999, *ApJ*, 518, L95
- Titarchuk, L., Osherovich, V., & Kuznetsov, S. 1999, *ApJ*, 525, L129
- van der Klis, M. 1995, in *X-ray Binaries*, ed. W.H.G. Lewin, J. van Paradijs, & E.P.J. van den Heuvel (Cambridge: Cambridge




**Fig. 4.** The rms amplitude versus the photon energy for NBOs in Scorpius X-1. The solid lines are the best fits with a power law to the data points in the five lower energy bands.

Univ. Press), 252

van der Klis, M. 1989, in Timing neutron stars, eds. Ögelman, H., & van den Heuvel, E. P. J. NATO Advanced Science Institutes (ASI) Series C, 262, p.27

van der Klis, M., Swank, J. H., Zhang, W., et al. 1996, ApJ, 469, L1

van der Klis, M., Wijnands, R. A. D., Horne, K., & Chen, W. 1997, ApJ, 481, L97

Yu, W. 2007, ApJ, 659, L145

Yu, W., van der Klis, M., & Jonker, P. G. 2001, ApJ, 559, L29